iFluor® 430 Goat Anti-human IgG (H+L) Antibody *Cross Adsorbed*

Catalog Number: 50050, 50051
Unit Size: 200 ug, 1 mg

Product Details

Storage Conditions	2-8°C with minimized light exposure. Do not freeze.
Expiration Date	12 months upon receiving
Concentration	1 mg/mL
Formulation	Phosphate-buffered saline (PBS, pH 7.2), 2 mg/mL BSA

Unit Details

Reconstitution Volume	**50050 (200 ug)**	**50051 (1 mg)**
	200 uL ddH_2O	1 mL ddH_2O

Antibody Properties

Species Reactivity	Human
Class	Secondary
Clonality	Polyclonal
Host	Goat

Biological Properties

Stabilizer	2 mg/mL BSA
Appearance	Solid
Preparation	Goat anti-human IgG (H+L) is produced in goat with pooled total human IgG and affinity purified with human IgG coupled beads. The antibody is conjugated with iFluor® 430 under optimal conditions.
Application	Flow Cytometry (FACS), IF, IHC, ELISA, WB
Recommended Dilutions	Suggested dilutions are only guidelines; users should titrate the product for their specific assay using appropriate controls

Application	Recommended dilution
Flow Cytometry (FACS)	1-5 µg/mL

©2025 AAT Bioquest, Inc. Last revised December 2025. For more information and tools, please visit https://www.aatbio.com

IF	2 µg/mL
IHC	1-10 µg/mL
ELISA	100 ng/mL
WB	1-10 µg/mL

Spectral Properties

Conjugate	iFluor™ 430
Excitation Wavelength	433 nm
Emission Wavelength	498 nm

Applications

AAT Bioquest's anti-human secondary antibodies have well-characterized specificity for human immunoglobulins and are useful in the detection, sorting or purification of its specified target. This iFluor® 430-labeled secondary antibody was prepared using AAT Bioquest's proprietary labeling technology. It demonstrated much brighter signal compared to the similar iFluor® 430 goat anti-human IgG antibodies from other commercial sources, and thus can significantly increase assay sensitivities. Secondary antibodies offer increased versatility enabling users to use many detection systems (e.g. HRP, AP, fluorescence). They can also provide greater sensitivity through signal amplification as multiple secondary antibodies can bind to a single primary antibody. This antibody was purified through affinity chromatography and conjugated to iFluor® 430 (ex/em = 433/498 nm). It is compatible with the 445 nm laser and 510/80 nm bandpass filter (for example, as in the BD FACSAria™ III).

©2025 AAT Bioquest, Inc. Last revised December 2025. For more information and tools, please visit https://www.aatbio.com